

<https://helda.helsinki.fi>

Sydnone anions and abnormal N-heterocyclic carbenes of O-ethylsydnones : Characterizations, calculations and catalyses

Wiechmann, Sascha

2014

Wiechmann , S , Freese , T , Drafz , M H H , Hübner , E G , Namyslo , J C , Nieger , M & Schmidt , A 2014 , ' Sydnone anions and abnormal N-heterocyclic carbenes of O-ethylsydnones : Characterizations, calculations and catalyses ' , Chemical Communications , vol. 50 , no. 80 , pp. 11822-11824 . <https://doi.org/10.1039/c4cc05461j>

<http://hdl.handle.net/10138/224668>

<https://doi.org/10.1039/c4cc05461j>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.



Cite this: *Chem. Commun.*, 2014, 50, 11822

Received 15th July 2014,
Accepted 15th August 2014

DOI: 10.1039/c4cc05461j

www.rsc.org/chemcomm


Sydnone anions and abnormal N-heterocyclic carbenes of O-ethylsydnones. Characterizations, calculations and catalyses†

Sascha Wiechmann,^a Tyll Freese,^a Martin H. H. Drafz,^a Eike G. Hübner,^a
Jan C. Namyslo,^a Martin Nieger^b and Andreas Schmidt^{*a}

Deprotonated sydnones, which can be represented as anionic N-heterocyclic carbenes, were prepared as Li adducts and compared with deprotonated O-ethylsydnones (5-ethoxy-1,2,3-oxadiazol-4-ylidenes) which belong to the class of abnormal NHCs. The Pd complexes of the sydnone anions (X-ray analysis) as well as of the O-ethylsydnone carbenes proved to be efficient catalysts in aryl couplings of thiophenes.

Of all the 228 theoretically predicted mesoionic compounds¹ sydnones are probably the most prominent. Sydnones (1,2,3-oxadiazolium-5-olates) are 5-membered representatives of the class of conjugated mesomeric betaines (CMB) which is one of four distinct categories of this class of compounds.² They are not only known as versatile 1,3-dipoles in [2+3]-cycloadditions but also as biologically active compounds.³ So far the chemistry of sydnones has not been associated with the chemistry and applications of N-heterocyclic carbenes which have developed rapidly during the last two decades. Numerous structural variations of N-heterocyclic carbenes have been described, most of them aiming at enhancing the electron density at the carbene center and thus influencing the σ -donor capacity in catalytically active metal complexes.⁴ Recently mesomeric betaines including mesoionic compounds, however, have also come into the focus of carbene research. First results of interconversions of mesomeric betaines into N-heterocyclic carbenes and *vice versa* are summarized in recent review articles.⁵ In continuation of our interest in N-heterocyclic carbenes and mesomeric betaines⁶ we report here on sydnones as well as on O-ethylsydnones (5-ethoxy-1,2,3-oxadiazolium salts) and deprotonated species derived thereof.

We prepared a series of sydnones **2a–f** by cyclization of N-nitroso amino acids **1a–f** with acetic anhydride (Scheme 1). Among eight



Scheme 1

dipolar resonance forms of sydnones, **I** is the most common. The best single representation which reflects the properties of sydnones, however, appears to be **II**.⁷ Resonance form **II** is in agreement with carbonyl stretching frequencies of **2a–f** between 1728 cm^{−1} and 1736 cm^{−1}. Bond length (1.197 Å), calculated from IR spectra of **2a**, and bond order (1.99) correspond to an exocyclic C=O double bond which can also be seen in a single crystal X-ray analysis of **2b** (ESI†). We calculated the NICS(1)⁸ value of **2a** (−6.8) which is between the cyclopentadienyl anion (−12.3) and cyclopentane (−2) (DFT LACVP*/B3LYP). In summary these values confirm that sydnones are no aromatics.

In contrast to other mesoionic compounds such as nitron **3**⁹ and imidazolium-4-aminide **4**¹⁰ no hints on tautomeric equilibria between sydnones and their N-heterocyclic carbenes can be found in the spectra. DFT calculations revealed that the mesoionic form **2aA** is considerably more stable than its tautomer **2aB** [$\Delta E = +191$ kJ mol^{−1}] (Scheme 2). Except for few photochromic sydnones,¹¹ valence isomers such as **2aC** are not detectable.



A base screening employing LiOtBu, NaOtBu, KOtBu, KH, LiAlH₄ and *n*BuLi in a variety of solvents, respectively, which have been used earlier to deprotonate sydnones for further functionalizations at C4,^{3,7} caused a decomposition of the sydnones **2a–f**, when we tried to isolate the deprotonated species. Fortunately, deprotonation by cyanomethylolithium, freshly prepared from *n*BuLi and anhydrous MeCN, resulted in the formation of the sydnone anions **5a–f** in quantitative yields

^a Clausthal University of Technology, Institute of Organic Chemistry, Leibnizstrasse 6, D-38678 Clausthal-Zellerfeld, Germany. E-mail: schmidt@ioc.tu-clausthal.de

^b Laboratory of Inorganic Chemistry, Department of Chemistry, University of Helsinki, P.O. Box 55 (A.I. Virtasen aukio 1), FIN-00014, Helsinki, Finland

† Electronic supplementary information (ESI) available: Crystallographic data of **2b** and *trans*-**16**. CCDC 987516 and 987969. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4cc05461j





as colorless, moisture-sensitive compounds which proved to be stable under an inert atmosphere for several weeks (Scheme 3). On exposure to water or protic solvents, reprotonation of **5a-f** to the sydrones **2a-f** occurred, which were recovered quantitatively in all cases. Ethylation of the sydrones **2a-f** by triethyloxonium tetrafluoroborate gave the 5-ethoxy-1,2,3-oxadiazolium salts **6a-f** which are stable under an inert atmosphere up to approximately 60 °C. As the calculated NICS(1) value of **6a** is -8.9 these *O*-ethylsydrones are slightly more aromatic than the corresponding sydrones. The isolation of the abnormal N-heterocyclic carbenes (aNHC) **7a-f**, however, failed, as all attempts to deprotonate the *O*-ethylsydrones resulted in decomposition products.

Sydnone anions can be represented by several resonance forms, among those representations as anionic abnormal NHCs **I** and **II**, and the anionic NHC **III**. Do sydnone anions bear more than a formal relationship to N-heterocyclic carbenes? As a matter of fact, the calculated bond lengths of C4–C5 (1.451 Å), C5=O (1.227 Å), and N3–C4 (1.353 Å) are best represented by resonance form **II**. As calculated by us and found IR-spectroscopically,¹² the lithium cation is located between C4 and the exocyclic oxygen of **5a**. By contrast, the aNHC **7a** is best represented as shown, as the calculated C4–C5 bond (1.390 Å) is shortened in comparison to **5a**, whereas the C5–O and O–Et bonds are essentially $\text{C}_{\text{sp}^2}\text{--O}$ and $\text{O--C}_{\text{sp}^3}$ single bonds (1.325 and 1.445 Å, resp.). The ESP partial charges of C4 of the sydnone anion **5a** as well as of the aNHC **7a** adopt values between the CAAC **9**¹³ on the one hand, and the aNHC **10**¹⁴ on the other (Fig. 1). As a comparison, we also calculated the phenyl derivatives of carbene **11**¹⁵ derived from a cross-conjugated mesomeric betaine, the anionic NHCs **12**¹⁶ and **14**¹⁷ derived from the mesoionic compounds imidazolium-4-aminide and




Fig. 1 Calculated partial charges of the title compounds and selected NHCs.





Fig. 2 Calculated highest occupied molecular orbitals.


imidazolium-4-olate, respectively, and the NHC imidazole-2-ylidene **13**. Similar to N-heterocyclic carbenes derived from ylides¹⁸ the calculated HOMO of **5a** is a π -orbital with a large coefficient on C4, whereas the HOMO – 1 is perpendicular to the HOMO and contains the lone pair of electrons at C4 (Fig. 2).

As expected the HOMO of the aNHC **7a** is mainly a σ -type molecular orbital as shown.

Palladium complexes to stabilize the aNHC **7a** and to study catalytic activities were prepared as follows. 4-Bromo-*N*-phenylsydnone **15** formed the palladium complex **16** of the sydnone anion **5a** as a mixture of *trans* and *cis* isomers (Scheme 4).¹⁹ We were able to perform a single crystal X-ray analysis of *trans*-**16** (ESI[†]). Ethylation of the bromosydnone **15** under exclusion of moisture resulted in the formation of the new 4-bromo-1,2,3-oxadiazolium salt **17** in good yields, which served as starting material for the preparation of the Pd complex **18** of the aNHC **7a**.

The Pd complexes **16** and **18** of the sydnone anion **5a** as well as the aNHC **7a** proved to be efficient catalysts in Suzuki–Miyaura





Scheme 5

Table 1 Catalysts and yields

Entry	Starting material	Product	Catalyst	Yield ^a (%)
1	19a	20a	16	55
2	19a	20a	18	74
3	19a	20b	16	53
4	19a	20b	18	60
5	19b	20c	16	91
6	19b	20c	18	85
7	19b	20d	16	80
8	19b	20d	18	79

^a Isolated yields.

cross-coupling reactions (Scheme 5). Thus, the perphenylation of tetrabromothiophene **19a** to **20a** proceeds in considerably higher yields and in shorter periods of time under the conditions shown (Table 1, entries 1 and 2) in comparison to the catalyst system $\text{Pd(PPh}_3)_4/\text{K}_3\text{PO}_4$ (tol, H_2O , 12 h, 90°C) which gave 37% yield of **20a**.²⁰ The pernaphtylation of **19a** to the hitherto undescribed 2,3,4,5-tetra(naphthalen-1-yl)thiophene **20b** gave 53% and 60% yield, respectively, when the complexes **16** and **18** were employed (entries 3 and 4). The 2,5-bisarylation of 2,5-dibromo-3,4-dinitrothiophene **19b** with phenylboronic acid gave 3,4-dinitro-2,5-diphenylthiophene **20c** in excellent yields under catalysis of **16** and **18**, respectively (entries 5 and 6). Compound **20c** has been prepared before in 37% yield by cross-coupling using $\text{Pd(PPh}_3)_4$, K_2CO_3 in MeOH and toluene over a period of 8 h at reflux temperature.²¹ An alternative procedure uses 1,2-dimethoxyethane as solvent to give 80%, however, after a reaction time of 13 h at reflux temperature.²² Whereas the cross-coupling to 2,5-di(naphthalen-1-yl)-3,4-dinitrothiophene **20d**, which is a new compound, under $\text{Pd(PPh}_3)_4$ catalysis proceeded in only 40% yield, catalysis by **16** or **18** gave very good yields (entries 7 and 8).

Sydnone anions share typical features with N-heterocyclic carbenes, and their deprotonated O-ethyl derivatives, 5-ethoxy-1,3,4-oxadiazol-4-ylidenes, belong to the class of abnormal NHCs. Either species form stable Pd complexes which are efficient catalysts in Suzuki–Miyaura cross-coupling reactions.

Notes and references

- W. D. Ollis and C. A. Ramsden, *Adv. Heterocycl. Chem.*, 1976, **19**, 1; C. A. Ramsden, in *Comprehensive Organic Chemistry*, ed. P. G. Sammes (Hrsg.), Pergamon Press, Oxford, 1979, vol. 4, p. 1171.
- W. D. Ollis, S. P. Stanforth and C. A. Ramsden, *Tetrahedron*, 1985, **41**, 2239.
- D. L. Browne and J. P. Harrity, *Tetrahedron*, 2010, **66**, 553; T. L. Gilchrist, *Sci. Synth.*, 2004, **13**, 109; R. Chandrasekhar and M. J. Nanjan, *Mini-Rev. Med. Chem.*, 2012, **12**, 1359.
- Selected publications: M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485; J. C. Bernhammer, N.-X. Chong, R. Jothibas, B. Zhou and H. V. Huynh, *Organometallics*, 2014, **33**, 3607; Q. W. Zhao, D. P. Curran, M. Malacria, L. Fensterbank, J. P. Goddard and E. Lacôte, *Chem. – Eur. J.*, 2011, **17**, 9911; J. A. Mata and M. Poyatos, *Curr. Org. Chem.*, 2011, **15**, 3309; *N-Heterocyclic carbenes, from laboratory curiosities to efficient synthetic tools*, ed. S. Díez-González, Royal Society of Chemistry Publishing, Cambridge, 2011; T. Dröge and F. Glorius, *Angew. Chem., Int. Ed.*, 2010, **49**, 6940; G. Guisado-Barrios, J. Bouffard, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 4759; A. Schmidt, A. Beutler, M. Albrecht and F. J. Ramírez, *Org. Biomol. Chem.*, 2008, **6**, 287; F. E. Hahn and M. C. Jahnke, *Angew. Chem., Int. Ed.*, 2008, **47**, 3122; D. Holschumacher, T. Bannenberg, C. G. Hrib, P. G. Jones and M. Tamm, *Angew. Chem., Int. Ed.*, 2008, **47**, 7428; V. Lavallo, Y. Canac, B. Donnadiou, W. W. Schöller and G. Bertrand, *Angew. Chem., Int. Ed.*, 2006, **45**, 3488; A. J. Arduengo III, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361.
- A. Schmidt, S. Wiechmann and T. Freese, *ARKIVOC*, 2013, **i**, 424; A. Schmidt and Z. Guan, *Synthesis*, 2012, 3251; A. Schmidt and A. Dreger, *Curr. Org. Chem.*, 2011, **15**, 2897; A. Schmidt and A. Dreger, *Curr. Org. Chem.*, 2011, **15**, 1423; A. Schmidt, A. Beutler and B. Snovydyovych, *Eur. J. Org. Chem.*, 2008, 4073.
- Selected publications: Z. Guan, S. Wiechmann, M. Drafz, E. Hübner and A. Schmidt, *Org. Biomol. Chem.*, 2013, **11**, 3558; A. Schmidt and A. Rahimi, *Chem. Commun.*, 2010, **46**, 2995; A. Schmidt, N. Münster and A. Dreger, *Angew. Chem., Int. Ed.*, 2010, **49**, 2790; A. Schmidt and B. Snovydyovych, *Synthesis*, 2008, 2798; A. Schmidt and T. Mordhorst, *Synthesis*, 2005, 781.
- T. L. Gilchrist and P. M. O'Neill, *Comprehensive Heterocycl. Chem. II*, 1996, p. 165.
- M. Zora and I. Özkan, *THEOCHEM*, 2003, **638**, 157.
- C. Färber, M. Leibold, C. Bruhn, M. Maurer and U. Siemeling, *Chem. Commun.*, 2012, **48**, 227.
- V. César, J.-C. Tourneux, N. Vujkovic, R. Brousses, N. Lugan and G. Lavigne, *Chem. Commun.*, 2012, **48**, 2349; A. A. Danopoulos, K. Yu. Monakhov and P. Braunstein, *Chem. – Eur. J.*, 2013, **19**, 450.
- S. Nešpůrek, J. Lukáš, S. Böhm and Z. Bastl, *J. Photochem. Photobiol., A*, 1994, **84**, 257; S. Nešpůrek, S. Böhm and J. Kuthan, *THEOCHEM*, 1986, **136**, 261.
- M. G. Ezernitskaya, B. V. Lokshin, E. I. Kazimirschuk, V. N. Khandozhko and V. N. Kalinin, *Mikrochim. Acta, Suppl.*, 1997, **14**, 381.
- V. Lavallo, Y. Canac, C. Präsang, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2005, **44**, 5705.
- R. Tonner, G. Heydenrych and G. Frenking, *ChemPhysChem*, 2008, **9**, 1474.
- G. Lavigne, V. César and N. Lugan, *Chem. – Eur. J.*, 2010, **16**, 11432; V. César, N. Lugan and G. Lavigne, *J. Am. Chem. Soc.*, 2008, **130**, 11286.
- A. A. Danopoulos and P. Braunstein, *Chem. Commun.*, 2014, **50**, 3055.
- L. Benhamou, N. Vujkovic, V. César, H. Gornitzka, N. Lugan and G. Lavigne, *Organometallics*, 2010, **29**, 2616; L. Benhamou, V. César, H. Gornitzka, N. Lugan and G. Lavigne, *Chem. Commun.*, 2009, 4720; A. T. Bijou, K. Hirano, R. Fröhlich and F. Glorius, *Chem. – Asian J.*, 2009, **4**, 1786.
- N. Pidlypnyi, J. C. Namyslo, M. H. H. Drafz, M. Nieger and A. Schmidt, *J. Org. Chem.*, 2013, **78**, 1070.
- Some Pd complexes have been described before: V. N. Kalinin, F. M. She, V. N. Khandozhko and P. V. Petrovskii, *Russ. Chem. Bull.*, 2001, **50**, 525; V. Kalinin, S. Fan Min and P. Petrovskii, *J. Organomet. Chem.*, 1989, **379**, 195.
- T. T. Dang, N. Rasool, T. T. Dang, H. Reinke and P. Langer, *Tetrahedron Lett.*, 2007, **48**, 845.
- Q. Li, J. Li, R. Yang, L. Deng, Z. Gao and D. Liu, *Dyes Pigm.*, 2012, **92**, 674.
- H. Cheng, B. Djukic, H. A. Jenkins, S. I. Gorelsky and M. T. Lemaire, *Can. J. Chem.*, 2010, **88**, 954.

